Lizzy Compton and Terra Ficke

Kouma-Kondara Monthly Min data

In[1]:=

```
KoumaKondaraMinTemp =
  Flatten[Import["C:\\MATH375\\MinProject2\\Min_Temps_Kouma.xlsx"], 4];
MinYears = Range[KoumaKondaraMinTemp[[2, 1]], KoumaKondaraMinTemp[[-2, 1]]];
MinTemps = KoumaKondaraMinTemp[[2 ;; -2, 2]];
MinData = Table[{MinYears[[i]], MinTemps[[i]]}, {i, Length[MinYears]}];
lm = LinearModelFit[MinData, {x, Sin[2*Pi*x], Cos[2*Pi*x]}, x]
```

Out[2]= FittedModel[-41.32+0.0307556x+0.668216 Sin[2 Pi x]+0.644844 Cos[2 Pi x]]

In[3]:=

```
(*I don't know why my intercept is at -41..*)
p1 = ListPlot[MinData];
p2 = Plot[lm[x], {x, 1961, 2016}];
Show[p1, p2]
```

Out[8]=



Out[10]=



In[11]:=
```
linearModel["RSquared"]
```
Out[11]=
```
0.468
```

In[12]:=
```
linearModel["ParameterConfidenceIntervals"]
```
Out[12]=
```
{{-48.1188, -34.5223}, {0.0273371, 0.0341574},
 {0.567782, 0.721496}, {0.00848698, 0.0145931}}
```

In[13]:=
```
(*The linear model is significant all parameters except the intercept*)
```

In[14]:=
```
ListPlot[linearModel["FitResiduals"], Filling -> Axis,
 PlotLabel -> Hold[Residuals vs Time], LabelStyle -> {GrayLevel[0]}]
```
Out[14]=



In[15]:=
```
qmLinearModel[MinDataTemp_, Sin[2 Pi x], Cos[2 Pi x]}, x]
ShowListPlot[MinDataPlot[qm[x], {x, 1961, 2016}],
 PlotLabel -> Hold[Average Monthly Minimum Temperature vs Quadratic Model],
 LabelStyle -> {GrayLevel[0]}]
```
Out[15]=
```
FittedMo 5.71 - 0.0165 x + 0.0682 x^2 - 0.648 x]
```



In[16]:=
```
qm["RSquared"]
```
Out[16]=
```
0.468
```

In[18]:=
```
nlm[Parameter Confidence]
```
Out[18]=
```
{{-946.7, 958.1}, {-0.974, 0.941}}
{{-0.000228, 0.000902}, {-0.00854, 0.00597}}
```

In[19]:=
```
(*So the quadratic is significant except in the cos term*)
```

In[20]:=
```
ListPlot[nlm["FitResiduals"], PlotLabel -> "Residuals", AxesLabel -> {"Index", "Residual"}]
```
Out[20]=



In[21]:=
```
ListPlot[nlm["FitResiduals"], PlotLabel -> "Residuals", AxesLabel -> {"Index", "Residual"}, PlotStyle -> {GrayLevel[0.5]}]
```
Show 6 more lines

Out[21]=

In[22]:=

In[23]:=

Trying it again, removing anything below 17 or above 22 (I just picked those numbers to be the limit for the outliers because that seemed to make sense for the data)

Kouma-Kondata Monthly Min data

In[24]:=

In[25]:=
```
KoumaKondaMin2 = Partition[Flatten[Import["M:\\MATH375\\MinProject2\\Min_Temps.xlsx"]], 4];
```

In[26]:=
```
MinYearKouma2 = KoumaKondaMin2[[2, 1]];
MinTempKouma2 = KoumaKondaMin2[[2, 4]];
```

In[28]:=
```
MinDataTable = Table[{MinYearKouma2[[i]], MinTempKouma2[[i]]}, {i, Length[MinYearKouma2]}];
```

In[29]:=
```
lm2 = LinearModelFit[MinData, {x, Sin[2*Pi*x], Cos[2*Pi*x]}, x]
Show[ListPlot[MinData, PlotLabel -> Hold[AdjustedAverageMonthlyMinimumTemperatureQuadraticModel],
 LabelStyle -> GrayLevel[0]]
```

Out[29]=
```
FittedModel[ -40.8326 + 0.0303x + 19.9974 Sin[2 x] - 0.6358 Cos[2 x] ]
```



Out[30]=

In[31]:=

In[32]:=
```
ListPlot[lm2["FitResiduals"], Filling -> Axis,
 PlotLabel -> Hold[AdjustedResidualsTimeLabelStyle -> GrayLevel[0]]
```



Out[32]=

In[33]:=
```
(* There is still a pattern in the residuals *)
```

In[34]:=
```
lm2["RSquared"]
```

Out[34]=
```
0.573
```

In[35]:=
```
lm2["ParameterConfidenceIntervals"]
```

Out[35]=
```
{{-46.4293, -35.2359}, {0.0277055, 0.0333336}, {19.5722, 20.4226}, {-0.698114, -0.0343033}}
```

In[36]:= (*All parameters significant*)

In[39]:= qm = LinearModelFit[Table[{x, Sin[2 Pi x], Cos[2 Pi x]}, {x, 1961, 2016}], {1, x, Sin[2 Pi x], Cos[2 Pi x]}, x]
ListPlot[{MinDate[qm], Max[qm]}, {x, 1961, 2016}, Frame → True]

Out[39]= FittedModel[-642.2 + 0.3241 x + 0.0979 Sin[2 π x] + 0.6345 Cos[2 π x]]

Out[38]=

In[39]:= qm["RSquared"]

Out[39]= 0.5751

In[40]:= qm["ParameterConfidenceIntervals"]

Out[40]= {{-1424.3, 339.8}, {-0.1511, 0.4219}, {-0.0003499, 0.0004568}, {0.5716, 0.6974}}

In[41]:= (*Only the sin and cos parameters significant*)

In[42]:= ListPlot[qm["FitResiduals"], Filling → Axis]

Out[42]=